

PART I  
Molecular Clouds and  
Star Formation

DR.RUPNATHJI( DR.RUPAK NATH )

DR.RUPNATHJI( DR.RUPAK NATH )

We review recent results from numerical simulations and related models of MHD turbulence in the interstellar medium (ISM) and in molecular clouds. We discuss the implications of turbulence for the processes of cloud formation and evolution and the determination of clouds' physical properties. Numerical simulations of the turbulent ISM to date have included magnetic fields, self-gravity, parameterized heating and cooling, modeled star formation, and other turbulent inputs. The structures that form reproduce well the observed velocity-size scaling properties while predicting the nonexistence of a general density-size scaling law. Criteria for the formation of gravitationally bound structures by turbulent compression are summarized. For flows with equations of state  $P \propto \rho^\gamma$ , the statistics of the density field depend on the exponent  $\gamma$ . Numerical simulations of both forced and decaying MHD compressible turbulence have shown that the decay rate is comparable to the nonmagnetic case. For virialized clouds, the turbulent decay time is shorter than the gravitational free fall time, so wholesale cloud collapse is prevented only by ongoing turbulent inputs, a strong mean magnetic field, or both. Finally, perspectives for future work in this field are briefly discussed.

## I. INTRODUCTION

### A. Observational Motivation

Present-day star formation in our Galaxy is observed to take place in cold molecular clouds, which appear to be in a state of highly compressible magnetohydrodynamic (MHD) turbulence. Furthermore, the atomic gas

[3]

DR.RUPNATHJUK(DR.RUPAK NATH)

within which the molecular clouds form is also turbulent on larger scales, from hundreds to thousands of parsecs (e.g., Braun 1999). In this chapter we review recent results in the areas of cloud formation, structure, and evolution as well as their implications for observed physical and statistical cloud properties; these results are obtained mainly from numerical simulations of compressible MHD turbulence and from related analytical models. An introductory review, including a more detailed discussion of turbulence basics and discussions (current up to 1997) on fractality and phenomenological models, has recently been given by Vázquez-Semadeni (1999). The review of Ostriker (1997) focuses on the role of MHD turbulence in the internal dynamical evolution of molecular clouds. The reviews of Heiles et al. (1993) and McKee et al. (1993) provide extensive discussions of the observations and general theory, respectively, of magnetic fields in star-forming regions.

The formation of molecular clouds probably cannot be considered separately from the formation of larger diffuse H I structures, because the former are often observed to have H I “envelopes” (e.g., Peters and Bash 1987; Shaya and Federman 1987; Wannier et al. 1991; Blitz 1993; Williams et al. 1995). This suggests that molecular clouds and clumps may be regarded as the “tips of the icebergs” in the general continuum interstellar density field of galaxies. The process of cloud formation quite possibly involves more than a single mechanism, including the passage of spiral density waves and the effects of combined large-scale instabilities (e.g., Elmegreen 1993*a*, 1995; Gammie 1996), operating preferentially in the formation of the largest high-density structures, and the production of smaller density condensations by either swept-up shells (Elmegreen and Elmegreen 1978; Vishniac 1983, 1994; Hunter et al. 1986), or by a generally turbulent medium (Hunter 1979; Hunter and Fleck 1982; Tohline et al. 1987; Elmegreen 1993*b*). In Section II of this chapter we discuss recent results on the generation of density fluctuations in a turbulent medium, obtained from numerical simulations and semiphenomenological models. Other mechanisms of cloud formation have been reviewed extensively by Elmegreen (1993*a*, 1995). Note, however, that discrete cloud coagulation mechanisms discussed there may not be directly applicable in the context of a dynamically evolving continuum, such as that considered here.

Structurally, molecular clouds are very complex, with volume-averaged H<sub>2</sub> densities  $n(\text{H}_2)$  ranging from  $\lesssim 50 \text{ cm}^{-3}$ , for giant molecular clouds (GMCs) of sizes several tens of parsecs (e.g., Blitz 1993), to  $n(\text{H}_2) \gtrsim 10^5 \text{ cm}^{-3}$  for dense cores of sizes 0.03–0.1 pc (e.g., Wilson and Walmsley 1989; Plume et al. 1997; Pratap et al. 1997; Lada et al. 1997; Myers 1995). Kinetic temperatures are relatively constant, typically  $T \sim 10 \text{ K} \pm$  a few degrees (e.g., Pratap et al. 1997), although temperatures larger by factors of a few are found in the vicinity of star-forming regions (e.g., Torrelles et al. 1983; Solomon and Sanders 1985). Their internal velocity dispersions are generally supersonic, with Mach numbers up to  $\gtrsim 10$ , except for the smallest cores (size  $R \lesssim 0.1 \text{ pc}$ ) (e.g., Larson 1981; Blitz 1993 and ref-

ferences therein; Goodman et al. 1998). When available (and significant; see, e.g., Crutcher et al. 1993, 1996; Padoan and Nordlund 1999), Zeeman measurements give typical values of the magnetic field intensity of a few to a few tens of microgauss (e.g., Heiles et al. 1993; Crutcher et al. 1993; Troland et al. 1996; Crutcher 1999), consistent with near equipartition between the kinetic and magnetic energies.

Additionally, interstellar clouds seem to follow power law scaling laws between their average density, velocity dispersion, and size (Larson 1981; see also Blitz 1993 and references therein). Together with the near equipartition between kinetic and magnetic energy, these are normally interpreted as evidence for virialized magnetic support for the clouds (e.g., Shu et al. 1987; Mouschovias 1987; Myers and Goodman 1988*a,b*; see also Whitworth 1996), although they provide only arguments of self-consistency of the virial equilibrium hypothesis rather than conclusive proofs (Heiles et al. 1993). In addition to the clouds that are self-consistent with virialization, however, examples also exist of objects that appear to be highly disturbed (e.g., Carr 1987; Loren 1989; Plume et al. 1997), or simply regions away from map intensity maxima (Falgarone et al. 1992, 1998) that do not satisfy one or more of the scalings. Another example of such scalings is the mass spectrum of the clouds and clumps, which seems also to be a power law (e.g., Blitz 1993; Williams et al. 1995), although present galactic cloud identification surveys remain incomplete.

Molecular clouds also exhibit signatures of self-similarity or, more generally, multifractality. Scalo (1990) and Falgarone et al. (1991) have shown that cloud boundaries have projected fractal dimensions  $\sim 1.4$ . Also, clouds exhibit hierarchical structure; that is, the densest structures are nested within larger, less dense ones in a self-similar fashion (e.g., Scalo 1985), at least at large scales. The self-similarity appears either to break down or, at least to change similarity exponents at some small scale, reported by Larson (1995) to be close to 0.05 pc in Taurus. That scale corresponds to the local Jeans length (see section III.B of the chapter by Meyer et al., this volume), although the latter point is controversial (e.g., Simon 1997; Bate et al. 1998). A similar break has also been reported at scales 0.25–0.5 pc (Blitz and Williams 1997), showing that this question is still largely open.

Such complexity strongly suggests itself as a manifestation of the turbulent regime that permeates molecular clouds. Thus, an understanding of the basics of compressible MHD turbulence and its incarnation in the interstellar case is necessary for explaining molecular cloud structure and evolution and for diagnosing the consequences of turbulent dynamics for the process of star formation. The disordered, nonlinear nature of interstellar MHD turbulence makes direct numerical simulation the most useful tool for developing this understanding. In section III of this chapter, we summarize recent results on cloud structure and discuss questions such as whether the physical conditions and scaling laws observed in clouds arise naturally in simulations of the turbulent ISM, and what simulations

predict for energy balance in the clouds, density fluctuation statistics, and the general characterization of structure in these flows.

Evolutionary aspects of turbulence and the mechanisms of cloud support under highly nonlinear, strongly self-gravitating conditions have remained open questions since the original identification of turbulent molecular clouds in the interstellar medium (ISM). Maintenance of supersonic turbulence faces the well-known problem of an expected excessive dissipation in shocks (Goldreich and Kwan 1974). Arons and Max (1975) proposed that the “turbulent” motions in molecular clouds may actually be moderate-amplitude (sub-Alfvénic) MHD waves, arguing that shock formation and the associated dissipation would then be significantly diminished. Additionally, it has been suggested that even strong dissipation may be compensated by sufficient energy injection from embedded stars and other sources (e.g., Norman and Silk 1980; Scalo 1987; McKee 1989; Kornreich and Scalo 1999). Section IV in this chapter discusses recent numerical MHD results on calculating dissipation rates for turbulence with parameters appropriate for the cold ISM and the implications of these results for questions of cloud support against self-gravitating contraction. In section V we conclude with a summary and discussion of outstanding goals and challenges for future research.

## B. MHD Compressible Turbulence Basics

Before proceeding to the next sections, a few words on the nature and parameters of molecular cloud turbulence are in order. Its properties are very different from laboratory incompressible turbulence. Besides the viscous and magnetic Reynolds numbers, which measure the magnitude of the nonlinear advection term  $\mathbf{v} \cdot \nabla \mathbf{v}$  in the momentum equation as compared to the viscous and ohmic dissipation terms, additional nondimensional parameters are necessary to characterize the flow. Two of these parameters are the sonic (thermal) and Alfvénic Mach numbers  $M_s \equiv u/c$  and  $M_A \equiv u/v_A$ , where  $u$  is a characteristic velocity of the flow,  $c$  is the isothermal sound speed, and  $v_A = B/(4\pi\rho)^{1/2}$  is the Alfvén speed, with  $B$  the magnetic field strength and  $\rho$  the fluid density. The plasma beta,  $\beta \equiv c^2\rho/(B^2/8\pi) = 2(M_A/M_s)^2$  (i.e., the ratio of the thermal to the magnetic pressure), is also frequently used to characterize the importance of magnetic fields. The last basic parameter, which characterizes the importance of self-gravity, is the Jeans number  $n_J \equiv L/L_J = L/\sqrt{\pi c^2/G\rho}$ ; this compares the cloud linear scale  $L$  with the minimum Jeans-unstable wavelength  $L_J$ .

Other peculiarities of ISM turbulence are related to the driving mechanisms. In standard (“Kolmogorov”) incompressible turbulence theory, energy is assumed to be injected at large scales; from there it cascades down to the small scales where it is dissipated. In the ISM, on the other hand, forcing occurs at all scales (e.g., Scalo 1987; Norman and Ferrara 1996). Some mechanisms operate at large scales ( $\geq 1$  kpc), such as the galactic shear (Fleck 1981, although this is believed to be a very inefficient source;

see, e.g., Shu et al. 1987) and supershells. Others operate at intermediate scales ( $\geq 100$  pc), such as expanding H II regions and supernova explosions. Yet others act at small scales (a few tenths of a parsec to a few parsecs), such as stellar winds or bipolar outflows. All these mechanisms are important sources of kinetic energy.

Small-scale dissipation mechanisms in the cold ISM are of several types. First, there is the usual viscous dissipation, which operates mostly in shocks and which remains finite even in the limit of vanishing viscosity. A significant amount of dissipation also results from ambipolar diffusion due to ion-neutral friction at high densities in the presence of a strong enough magnetic field (Kulsrud and Pearce 1969; see McKee et al. 1993; Myers and Khersonsky 1995). Cooling processes are also very efficient and may radiate much of the dissipated energy, but they are balanced by heating processes and may result in near-isothermal or quasipolytropic behavior (i.e.,  $P \propto \rho^\gamma$ ) (section II).

Even aside from all these astrophysical properties, compressible turbulence has many distinctive features related to the transfer of energy from large scales to the dissipation (thermalization) scale. The new ingredient, as compared to incompressible turbulence, is the existence of *potential* (or compressible) modes in addition to the solenoidal (vortical) ones. Because compressions and rarefactions can exchange energy between bulk kinetic and microscopic thermal parts, compressible flows do not in general conserve bulk kinetic energy. Strongly radiative shocks, which yield an irreversible energy loss, are the most important aspect of this feature. The presence of additional compressive and thermal degrees of freedom therefore excludes the use of dimensional analysis to determine the slope of the velocity spectrum, as is done (by assumption of an energy-conservative cascade through scales) in the so-called K41 theory (Kolmogorov 1941; Obukhov 1941; see, e.g., Landau and Lifshitz 1987), which produces the well-known “universal” spectrum of the form  $d(u^2)/dk \equiv E(k) \propto k^{-5/3}$  for incompressible, nonmagnetic turbulence, where  $k = 2\pi/\lambda$  is the wavenumber corresponding to wavelength  $\lambda$ . For magnetized, incompressible, strong turbulence, recent theoretical work suggests that the mean magnetic field leads to strong anisotropy in the cascade but the same averaged energy spectrum (Goldreich and Sridhar 1995).

Moreover, the K41 theory is based on the assumption of locality in Fourier space of the nonlinear cascade (i.e., transfer between Fourier modes of similar wavelengths), a hypothesis probably not valid in the compressible case, because coupling among very different scales occurs in shocks. In these highly intermittent (i.e., inhomogeneous in space and time) structures, all Fourier modes decay at the same rate (Kadomtsev and Petviashvili 1973; Landau and Lifshitz 1987), thus invalidating the notion of inviscid cascade along the “inertial range,” defined as the range in Fourier space where the energy flux is constant. Note, however, that cascade processes in the presence of shocks have been discussed by

Kraichnan (1968) and Kornreich and Scalo (1999). The spectrum of the compressible modes appears to approach a Burgers (1974) spectrum of the form  $k^{-2}$ , which arises simply from the Fourier structure of the shocks. The corresponding configuration-space scaling for Burgers turbulence is  $u_l \propto l^{1/2}$  (section III.A). This scaling has been observed in numerical simulations of strongly compressible turbulence even in the presence of the magnetic field (Passot et al. 1988, 1995; Gammie and Ostriker 1996; Balsara et al. 1997; Stone 1998).

## II. TURBULENT CLOUD FORMATION

An important question is whether structures formed by either turbulent compressions or passages of single shock waves can become gravitationally unstable and collapse (Ögelman and Maran 1976; Elmegreen and Lada 1977; Elmegreen and Elmegreen 1978; Hunter 1979; Hunter and Fleck 1982; Hunter et al. 1986; Tohline et al. 1987; Stevens et al. 1999; Elmegreen 1993*b*; Klein and Woods 1998). As pointed out by Hunter et al. (1986), when both heating and cooling are present, the isothermal approximation, often used to describe radiative flows, is just one out of a continuum of possibilities. In cases where the heating and cooling rates are faster than the dynamical rates and are reasonably approximated by power law functions of the density and temperature, the gas can be described as a barotropic fluid with power law “equation of state”  $P \propto \rho^\gamma$  (e.g., de Jong et al. 1980; Maloney 1988; Elmegreen 1991; Vázquez-Semadeni et al. 1996); we use the term “effective polytropic exponent” to refer to  $\gamma$ .<sup>a</sup> In general,  $\gamma$  is different from 1 for the global ISM (Myers 1978) and possibly even for molecular clouds (de Jong et al. 1980; Scalo et al. 1998). Note, however, that, because cooling laws are often approximated by piecewise power laws, the effective polytropic exponent is also approximately piecewise constant. Numerical simulations that explicitly include heating and cooling appropriate for atomic gas indeed show that the corresponding rates are faster than the dynamical rates by factors of at least 50 (Passot et al. 1995; Vázquez-Semadeni et al. 1996), confirming earlier estimations (Spitzer and Savedoff 1950; Elmegreen 1993*b*). However, those simulations depart from pure power law equations of state near star formation sites. For optically thin gas with rapid heating and cooling, an effective polytropic equation of state with parameterized  $\gamma$  may therefore be considered as the next level of refinement over isothermal or adiabatic laws; from the aforementioned simulations,  $\gamma$  is found to take values between

<sup>a</sup> We remark that this effective polytropic exponent, which describes the thermal behavior of gases with local heating and cooling in equilibrium, is different from the adiabatic exponent  $c_p/c_v$  (describing thermal behavior with no heating or cooling) and that its usage does not imply in any form that the system is in hydrostatic equilibrium. Also note that, strictly speaking, the polytropic equation is not properly an equation of state, because the ideal-gas equation of state is also satisfied at all times.

0 and 1/2 in the warm and cold “phases.” For molecular material that is optically thick and turbulent, it is not yet clear from simulations how well an isothermal or effective polytropic law may be satisfied.

The choice of cooling and heating functions in the aforementioned simulations implies the absence of an isobaric thermal instability (Field et al. 1969) at densities  $\sim 0.5\text{--}5\text{ cm}^{-3}$  and temperatures between  $10^2$  and  $10^4$  K. There exist realistic values of interstellar parameters, such as the background UV field, coolant depletion, and grain abundances, for which such an instability is indeed not realized (e.g., Draine 1978; Wolfire et al. 1995). This choice of parameters has allowed the analysis of the role of turbulence separately from that of the thermal instability. Even though it is desirable to study these processes in combination, the thermal instability is unlikely to be the main driver of interstellar motions, because thermal pressure is smaller, by at least a factor of 3, than each of the other principal forms of interstellar pressure (turbulent, magnetic, and cosmic-ray), and at least  $\sim 10$  times smaller than their sum (e.g., Boulares and Cox 1990).

The stability of fluid parcels compressed in  $n$  dimensions by shocks or turbulence requires  $\gamma > \gamma_{\text{cr}} \equiv 2(1 - 1/n)$  (Chandrasekhar 1961; McKee et al. 1993; Vázquez-Semadeni et al. 1996). Shocks are likely to have  $n = 1$ , but generic turbulent compressions can have any dimensionality  $n \leq 3$ . The latter authors have also shown numerically the production of small-scale collapsing regions within a globally stable turbulent medium, in both magnetic and nonmagnetic cases, although the former still requires the collapsing region to be supercritical (cf. section IV.B). They also noted that turbulence-induced collapse generally involves small fractions of the mass available in the flow, in contrast to the case of globally unstable regions, and they showed that the density jump  $\rho_2/\rho_1 \equiv X$  across shocks in a polytropic medium satisfies  $X^{1+\gamma} - (1 + \gamma M_s^2)X + \gamma M_s^2 = 0$ . This equation recovers the well-known result that  $X = M_s^2$  for  $\gamma = 1$ , but it also implies that in the limit of vanishing  $\gamma$ ,  $X \rightarrow \exp(M_s^2)$ .

An important consequence of this effective polytropic behavior is that, if  $0 < \gamma < 1$ , denser regions are *colder*. Upon the production of turbulent density fluctuations, the flow develops a temperature distribution similar to that resulting from isobaric thermal instabilities ( $\gamma < 0$ ) but without the need for them. Essentially, turbulent ram pressure provides the drive that, in the thermally unstable case, is provided by thermal pressure. Note, however, that in the turbulent  $\gamma > 0$  case there are no sharp phase transitions. In this scenario, the near-constancy of the thermal pressure among the various moderate-density phases of the ISM is an incidental consequence of a small value of  $\gamma$  rather than a cloud-confining agent (Ballesteros-Paredes et al. 1998). The increase in thermal pressure in molecular clouds is a consequence of  $\gamma$  being closer to unity at the relevant densities (Scalo et al. 1998), although the thermal pressure is still subordinate to other sources of pressure for scales  $\geq 0.1$  pc.

For three-dimensional compressions, the minimum Mach number  $M_0$  necessary to induce collapse by the velocity field has been computed by

several authors as a function of  $\gamma$  and the mass  $m$  of the cloud in units of the Jeans mass. It is found that  $M_0 \propto \ln m$  for the isothermal ( $\gamma = 1$ ) case (Hunter 1979),  $M_0 \propto m^{-(\gamma-1)/(4-3\gamma)}$  for  $4/3 > \gamma > 1$  (Hunter and Fleck 1982), and  $M_0 \geq \sqrt{10/3}(1 - \gamma)$  for  $0 < \gamma < 1$  (Tohline et al. 1987). Note that the latter result is independent of the cloud's mass, at least for perfect spherical geometries.

Recently, Ballesteros-Paredes et al. (1999) have investigated some implications of the scenario that clouds are turbulent density fluctuations by exploring the properties of clouds formed in 2D numerical simulations of the ISM (Passot et al. 1995) on scales between 1.25 pc and 1 kpc, including standard atomic-gas cooling rates, diffuse background heating, modeled stellar ionization heating, self-gravity, the Coriolis force, galactic shear, and magnetic fields. It should be stressed that the clouds *form* during the evolution of the simulations, rather than being started with some pre-defined conditions, and they have estimated lifetimes (Ballesteros-Paredes et al. 1999) comparable to observational estimates (e.g., Bash et al. 1977; Blitz and Shu 1980; Blitz 1994). Ballesteros-Paredes et al. (1999) find that the velocity field is in general continuous across the boundaries, with kinetic surface terms in the Eulerian form of the virial theorem (McKee and Zweibel 1992) of comparable magnitude to the total kinetic energy contained within the clouds.

Ballesteros-Paredes et al. (1999) also suggest that the formation of hydrostatic structures within a turbulent medium is not possible unless the effective polytropic exponent  $\gamma$  increases during the process of collapse initiated by a turbulent compression. Such a change in  $\gamma$  will not occur until protostellar densities are reached if thermal pressure alone is considered, so in general the production of hydrostatic structures appears unlikely.

### III. CLOUD STRUCTURE

#### A. Scaling Relations and Energy Spectra

One of the most conspicuous properties of molecular clouds (shared with diffuse H I clouds), are the so-called Larson (1981) relations, which in their most common form read  $\Delta v \propto R^{1/2}$  and  $\rho \propto R^{-1}$ , where  $R$  is a characteristic spatial scale (e.g. cloud size),  $\Delta v$  is a characteristic velocity difference (e.g. linewidth or line centroid difference), and  $\rho$  the cloud's mean density.<sup>b</sup> As stated in the Introduction, the most generally accepted explanation of the origin of these relations is that the clouds are in virial equilibrium between gravity and turbulent support (which causes one of the two relations to become a consequence of the other); some other assumption, such as that the clouds are magnetically critical and that the magnetic field does not vary much from one region to another (e.g., Shu et al. 1987;

<sup>b</sup> Note that Larson's original exponents were slightly different, being 0.38 for the linewidth-size relation and  $-1.1$  for the density-size relation.

DR. RUPNATHI DR. RUPAK NATHI

Mouschovias 1987), fixes the other relation. However, molecular clouds may not always be gravitationally bound (Blitz 1994); indeed, except at the largest scales, observed internal clumps within clouds are *not* gravitationally confined (Bertoldi and McKee 1992). Being highly dynamic entities, even bound clouds may generally not be in *static* virial equilibrium (Ballesteros-Paredes and Vázquez-Semadeni 1997).

Alternative explanations under turbulent conditions have been proposed as well. Kolmogorov-like arguments based on, for example, cascades of angular momentum (Henriksen and Turner 1984) or kinetic energy density (Ferrini et al. 1983; Fleck 1996), are reviewed in Vázquez-Semadeni (1999). Here we briefly discuss results from numerical simulations.

As mentioned in Section I.B, for highly compressible regimes the turbulent energy spectrum is expected to approach the form  $E(k) \propto k^{-2}$  of shock-dominated Burgers turbulence. If we assume that the observed linewidths measure the root-mean-square turbulent velocity  $u_l$  over scales smaller than  $l$ , we can write  $u_l^2 = 2 \int_{2\pi/l}^{\infty} E(k) dk \propto \int_{2\pi/l}^{\infty} k^{-2} dk \propto l$ , so that the observed velocity dispersion–size relation appears to emerge naturally. However, the identification of  $\Delta v(l)$  with  $u_l$  is not trivial. While  $\Delta v(l)$  is the linewidth within a beam of width  $l$  integrated over the whole line of sight,  $u_l$  is an average over an idealized ensemble (or, in practice, over all space) for volumes of size  $l^3$ . Thus, the identification of  $u_l$  with  $\Delta v(l)$  may depend on each beam being dominated by a single component of scale  $\sim l^3$ ; hierarchical density fluctuations could potentially produce the required structure. The data of Issa et al. (1990) and Falgarone et al. (1992), which include positions in the sky away from brightness maxima and still exhibit a similar relation (slope  $\sim 0.4$ ), seem to support the turbulent origin of the  $\Delta v$ - $R$  relationship (Issa et al. also considered random positions in their maps). Peng et al. (1998) have recently studied a sample of CCS clumps, most of which are reported to be gravitationally unbound yet seem to follow Larson’s linewidth-size relation as well. However, Bertoldi and McKee (1992) found that the smaller, non-self-gravitating clumps in their large study sample did not follow Larson’s linewidth-size relation.

Vázquez-Semadeni et al. (1997) have surveyed the clouds that appear in the 2D numerical simulations of Passot et al. (1995), finding a velocity dispersion-to-size relation with a logarithmic slope  $\sim 0.4$  as well (albeit with a large scatter). The energy spectrum of those simulations is indeed of the form  $k^{-2}$ . To study the development of the energy spectrum, simulations of MHD turbulence in which driving is localized in wavenumber space at scales smaller than the box have been performed in slab geometry (“ $1\frac{2}{3}$ D”) (Gammie and Ostriker 1996) and in fully 3D geometry (Stone 1999; Stone et al. 1998). For both cases, extended spectra develop in  $k$ -space above and below the range of driving frequencies, indicating the development of both “direct” and “inverse” cascades. The  $1\frac{2}{3}$ D models, which were evolved over very long times, show spectral slopes  $-2$  or slightly steeper both above and below the forcing scale.

The 3D models, although more limited in dynamic range, show combined  $u_k^2 + B_k^2$  slopes between  $-2$  and  $-\frac{5}{3}$  for  $k$  larger than the forcing scale, with the more negative values occurring in weaker magnetic field models. The slope of  $u_k^2$  alone is  $-2$  (stronger fields) or slightly steeper (weaker fields).

Overall, turbulent MHD simulations tend to evolve toward global energy equipartition. For example, Ostriker et al. (1998) and Stone et al. (1998) find perturbed magnetic energies between 30 and 60% of kinetic energies over a wide range of  $\beta$ . Kinetic and magnetic spectra are within factors of a few from each other (Passot et al. 1995), implying equipartition at all scales.

Vázquez-Semadeni et al. (1997) also find cloud mass spectra of the form  $dN(M)/dM \propto M^{-1.44 \pm 0.1}$ , consistent with the low end of observational estimates (e.g., Blitz 1993), suggesting that the simulations reproduce well a number of observational cloud properties, even though they are two-dimensional. Additionally, they find that Larson's density-size relation is not satisfied in general, but rather seems to be the upper envelope in a  $\log \rho$ - $\log R$  diagram of the clouds' locus. That is, it is satisfied only by the densest clouds at a given size (implying the largest column densities). Low-column-density clouds would naturally escape observational surveys that utilize limited amounts of integration time (Larson 1981; Kegel 1989; Scalo 1990), suggesting that this relation may be an observational artifact. The off-peak data of Falgarone et al. (1992) are consistent with this suggestion. In summary, these results can be interpreted as implying that a  $\Delta v$ - $R$  relation comparable to the observed scalings may be established globally as a consequence of the development of compressible MHD turbulence, while the density-size relation may occur only for gravitationally bound clouds within this turbulent field. Further work is necessary to confirm this possibility.

## B. Turbulent Pressure

Vázquez-Semadeni et al. (1998) have performed 2D and 3D numerical simulations of isothermal gravitational collapse in initially turbulent clouds, following the evolution of the velocity dispersion as the mean density increases during the collapse. They found power law behavior of the form  $P_t \sim \rho^{\gamma_t}$  for the "turbulent pressure." In particular, for slowly collapsing magnetic simulations,  $\gamma_t \sim 3/2$ , consistent with the result of McKee and Zweibel (1995) for the adiabatic exponent of Alfvén waves upon slow compression. However, nonmagnetic and rapidly collapsing (shorter Jeans length) magnetic simulations have  $\gamma_t \sim 2$ . Gammie and Ostriker (1996) also verified the McKee and Zweibel scalings  $P_{\text{wave}} \propto \rho^{3/2}$  and  $P_{\text{wave}} \propto \rho^{1/2}$  for Alfvén wave pressure of, respectively, an adiabatically contracting medium and propagation along a density gradient. For freely evolving decay simulations, however, they found only a weak, variable correlation between perturbed magnetic pressure and density. These results are incompatible with the so-called "logatropic" equation  $P \sim \ln \rho$ , warning against its usage in dynamical situations.

DR. RUPAK NATH

### C. The Density Field

The turbulent density field has a number of relevant statistical and physical properties. Besides the density-size scaling relation (Section III.A above), which may or may not be a true property of clouds, these exhibit hierarchical nesting (e.g., Scalo 1985; Houllahan and Scalo 1990, 1992; Vázquez-Semadeni 1994) and evidence for fractal (e.g., Falgarone et al. 1991) or even multifractal structure (Chappell and Scalo 1998). The spatial and statistical distribution of the density field is crucial in the study of star formation and the understanding of the stellar initial mass function (IMF).

To construct a real theory of the IMF (or at least of the cloud mass spectrum, if the actual masses of stars turn out to be rather independent of their parent clumps), it is necessary to have a complete knowledge of the density statistics. A simple theory based exclusively on the probability density function (PDF, also commonly referred to as the density distribution function) of the fluid density (Padoan 1995; Padoan et al. 1997*b*) has been criticized by Scalo et al. (1998), who point out that in addition to the probability of occurrence of high-density sites, it is also necessary to know how much mass is contained in these fluctuations; this requires information on multipoint statistics.

As a first step toward the understanding of density fluctuations in the ISM, the PDF of the density in polytropic gas dynamics ( $P \propto \rho^\gamma$ ) has been investigated as a function of the rms Mach number,  $\gamma$ , and the mean magnetic field strength (Vázquez-Semadeni 1994; Scalo et al. 1998; Passot and Vázquez-Semadeni 1998; Nordlund and Padoan 1999; Ostriker et al. 1999; Stone et al. 1998). In the isothermal case ( $\gamma = 1$ ) the density PDF is close to a lognormal distribution for every value of the Mach number. For polytropic cases with  $\gamma < 1$  or  $\gamma > 1$ , a power law develops at densities larger than the mean or smaller than the mean, respectively, the effect being enhanced as the Mach number increases. This behavior is a consequence of the dependence of the local Mach number on the density in each case (Passot and Vázquez-Semadeni 1998; Nordlund and Padoan 1998). It has been verified in one-dimensional numerical simulations of a forced polytropic gas, but it also appears to be supported in several dimensions and in the presence of thermal heating and cooling, which yield “effective” polytropic behavior (Scalo et al. 1998). In the limit of high Mach numbers or vanishing effective polytropic exponent, the behavior does not coincide with that of the Burgers equation (Passot and Vázquez-Semadeni 1998). One implication of these results is that it should be possible in principle to determine the actual effective  $\gamma$  of the medium from observational determinations of its PDF, provided the problem of deconvolving the projected PDF is solved and  $\gamma$  does not vary much in the observed region. Also, for example, the observation by Scalo et al. (1998) that full simulations of the ISM that include the turbulent magnetic field have PDF consistent with  $\gamma < 1$  suggests that the MHD waves do not give appreciably large values of  $\gamma$  (as is also suggested by the  $P \propto \rho^{1/2}$  scaling of propagating linear-amplitude waves; cf. McKee and Zweibel 1995).

The mean mass-averaged value of  $\log(\rho/\rho)$  increases with the Mach number  $M_s$ ; for isothermal models in 2.5D and 3D respectively, Ostriker et al. (1999) and Padoan et al. (1997a) both find a logarithmic dependence on  $M_s^2$ . In large-Mach-number, high-resolution 1D isothermal models by Passot and Vázquez-Semadeni (1998), a linear dependence is found between the variance of the density logarithm and  $M_s^2$ , leading to a mass-averaged value of  $\log \rho/\rho$  that varies like  $M_s^2$ ; this difference with higher-dimensional simulations (linear vs. logarithmic scaling with  $M_s^2$ ) likely arises because the 1D simulations have a purely compressive velocity field and the higher-dimension simulations do not. Stone et al. (1998) and Ostriker et al. (1999) show that the largest mean contrasts in the density logarithm occur in models with the strongest mean magnetic fields (see also Pouquet et al. 1990).

The spectrum of the density field has recently come under investigation. Padoan et al. (1997a) have reported a steep logarithmic slope,  $\sim -2.6 \pm 0.5$  for low-resolution 3D simulations of isothermal turbulence. Scalo et al. (1998) have reported two regimes, one with a slope  $\sim -1.9$  at low  $k$  and another with slope  $\sim -2.4$  at high  $k$  for high-resolution 3D simulations of the ISM with heating and cooling. The steeper slopes may be due to inadequate resolution. On the other hand, Lazarian (1995) has produced an algorithm for deconvolving projected HI interferometric spectra, favoring a slope  $\sim -1$ . Equivalent work for molecular clouds and high-resolution simulations in the corresponding regimes are necessary to resolve this issue.

An interesting application has been given by Padoan et al. (1997a), who have shown that a turbulent density field with a power law spectrum (though steep) and a lognormal PDF produces simulated plots of extinction dispersion vs. mean extinction that compare well with the analogous observational diagram for an extinction map of a dark cloud. Padoan and Nordlund (1999) also suggest that models with weaker mean magnetic fields (such that  $M_A \sim 10$ ) have extinction dispersion vs. mean extinction plots that show better agreement with observations than do models with stronger mean magnetic fields (such that  $M_A \sim 1$ ); however, these models do not include self-gravity, which would affect the distribution of column densities (i.e., extinctions).

#### D. Correlations among Variables

Numerical simulations are especially useful in the investigation of the spatial correlation among physical variables, a crucial ingredient in the formation of stars. Of particular interest is the correlation of magnetic field strength with density, as well as the correlation between field direction and the topology of density features.

The topology of the clouds is extremely clumpy and filamentary; for an example, see Color Plate 1. In general, the magnetic field exhibits a morphology indicative of significant distortion by the turbulent motions, with greater magnetic field tangling in cases with weak mean magnetic

fields (Ostriker et al. 1999; Stone et al. 1998). Magnetic fields may be either aligned with or perpendicular to density features (cf. Color Plate 1; Passot et al. 1995; Vázquez-Semadeni and Passot 1999; Ostriker et al. 1999; Stone et al. 1998), with the former trend most visible at the boundaries of supershells (Gazol and Passot 1998*a*), and the latter most prominent for strong-field simulations in which field kinks and density maxima coincide (Gammie and Ostriker 1996; Ostriker et al. 1999).

Globally, no clear trend between density and magnetic field intensity is found in the simulations, but very weak correlations are observed (Passot et al. 1995; Gammie and Ostriker 1996). For example, in the large-scale 2D ISM models of Passot et al. (1995), the field strength  $B$  varies from  $\sim 10^{-2} \mu\text{G}$  in the low-density intercloud medium to  $\sim 25 \mu\text{G}$  in the densest clouds ( $n \sim 50 \text{ cm}^{-3}$ , size  $\sim$  several tens of parsecs), although vanishing field strengths are also found in those regions. In 3D simulations with weak mean magnetic fields ( $B_0 = 1 \mu\text{G}$ ), Padoan and Nordlund (1999) found a large dispersion in the values of  $B$  as a function of density  $n$ , but a power law correlation  $B \propto n^{0.4}$  in the upper envelope of this distribution. In stronger-mean-field simulations ( $B_0 = 30 \mu\text{G}$ ), they found little variation of  $B$  with  $n$ . Padoan and Nordlund argue that the  $B$ - $n$  upper-envelope correlation found in the former case supports the notion that the mean fields in molecular clouds are weak, because a similar  $B$ - $n$  scaling has been observed for measured Zeeman field strengths (e.g., Crutcher 1998). However, because the range of densities over which the envelope correlation is found in simulations is much smaller than the density regime in which a power law  $B$ - $n$  relation is observed in real clouds, and because the simulations do not include gravity, the conclusion remains controversial.

To explore how line spectra may vary spatially in simulated clouds, Padoan et al. (1998) have produced synthetic non-LTE (not in local thermodynamic equilibrium) spectra of various molecular transitions from models with weak and strong mean magnetic fields. They find that their super-Alfvénic model reproduces the observational trend of linewidth vs. integrated temperature found by Heyer et al. (1996), whereas the equipartition model gives a weaker trend. However, these models do not include self-gravity, which would affect the density distributions and density-velocity correlations. Thus, it remains uncertain whether the weak-mean-field model of molecular clouds advanced by Padoan and Nordlund (1999) truly provides a better fit to observations.

## IV. CLOUD EVOLUTION

### A. Dissipation Rates and Turbulence Maintenance

As already mentioned, the first observations of supersonic internal cloud velocities immediately led to the questions, which have persisted up to the present, of what creates these large-amplitude motions and how they are

maintained. Since the large-scale ISM is itself turbulent, turbulent motions may be incorporated into cold clouds from their formation stages, being part of the same continuum. In addition, there may be ongoing inputs that tap energy from larger scales in the Galaxy or from smaller scales within the cloud, particularly associated with various aspects of star formation (see, e.g., Scalo 1987; Miesch and Bally 1994; Kornreich and Scalo 1999). One of the first steps in understanding cloud evolution and in relating this evolution to the initiation of star formation and dynamical feedback is to assess the rate of turbulent decay under the range of conditions representative of dark clouds and GMCs.

From dimensional analysis, the decay rate of kinetic energy per unit mass must scale as  $\dot{E} \sim v^3/R$ , where  $v$  is some characteristic speed (or weighted product of two or more speeds), and  $R$  is some characteristic scale. In incompressible turbulence (cf. section I.B), the only characteristic scales are that of the box ( $L$ ), that of the flow velocity difference  $u_L$  on the largest scale, and the value of the small-scale viscosity  $\nu$ . The velocity dispersion over the whole box  $\sigma_v \sim u_L$  provided the spectral slope is  $-1$  or steeper. The first two quantities determine the decay rate,  $\sigma_v^3/\nu$ , while the last sets the spatial dissipation scale to  $l_{\min}/L \sim (\nu/L\sigma_v)^{3/4} \equiv \text{Re}^{-3/4}$ . For a nearly pressureless (i.e., highly compressible) fluid, again possessing the same three characteristic scales, the same dissipation rate would apply, except that energy would be transferred by a shock directly from the largest scale to the dissipation scale within a flow-crossing time. For a magnetized flow of finite compressibility, on the other hand, other velocity scales in addition to  $\sigma_v$  (namely, those associated with thermal pressure and magnetic stress,  $c$  and  $v_A$ ) enter the problem and may potentially influence the scaling of the dissipation rate.

In molecular clouds the thermal pressure is very low, and therefore, strong dissipation in shocks is expected. However, from early observations until quite recently, it has widely been considered likely that the “cushioning” effect of magnetic fields would significantly reduce kinetic energy dissipation for motions transverse to the mean field, provided that turbulent velocities remain sub-Alfvénic. Low-dimension ( $1\frac{2}{3}$ D) simulations (Gammie and Ostriker 1996) provided some support for this idea in that they found a scaling of dissipation rate with  $\beta$  in quasisteady state as  $\dot{E} \propto \beta^{1/4}$ , such that magnetic fields of a few tens of microgauss could potentially provide a factor  $\sim 3$  reduction in dissipation compared to that in weak-field ( $\beta = 1$ ) cases. However, very recent higher-dimension numerical simulations of both forced and decaying turbulence have shown that although some differences remain between weak-field and strong-field cases, dissipation rates are never substantially lower than the predictions of unmagnetized turbulence. We next describe specific results.

Recent 3D MHD decay simulations (Mac Low et al. 1998; Stone et al. 1998; Padoan and Nordlund 1999) have followed the evolution of Mach-5 turbulence with a variety of initial velocity spectra and  $\beta$  ranging from

DR. PUPNATHI K. SURIYAK NATHI

0.02 to 2, and have also been compared with the results of unmagnetized models. These models have uniform initial  $\mathbf{B}$ . Stone et al. (1998) also included simulations of decay from fully saturated turbulence and from saturated turbulence with initial density fluctuations suppressed. These experiments all show kinetic energy decay times (defined as the time for kinetic energy reduction by 50%) in the range  $\sim 0.4$  to  $1t_f$ , where  $t_f$  is the flow-crossing time  $l/u_l$  on the main energy-containing scale. In Stone et al. (1998), the difference in decay time between the strongest-field case ( $\beta = 0.02$ , corresponding to  $44 \mu\text{G}$  for  $n_{\text{H}_2} = 10^3 \text{ cm}^{-3}$  and  $T = 10 \text{ K}$ ) and the unmagnetized run is less than a factor of two. In decay models, Mac Low et al. (1998) and Stone et al. (1998) find late-time power law dependence of the turbulent energy as  $E \propto t^{-\eta}$ , with  $\eta$  ranging between 0.8 and 1.

To assess turbulent decay rates for a quasisteady state, Stone et al. (1998) performed simulations in which a fixed mechanical power  $\dot{E}_K$  is input to the flow in the form of random, uncorrelated velocity perturbations. A saturated state with energy level  $E_K$  is reached after a time  $\sim t_f$ , with only relatively small differences in the saturation energy between magnetized and unmagnetized models; the turbulence dissipation times  $E_K/\dot{E}_K$  range between 0.5 and  $0.7 t_f$ . Mac Low (1999) has found similar results, and suggests that the power law time dependence of decay models arises from a secular increase in the smallest scale of turbulence. The implications of the short turbulence dissipation time for potential cloud support are discussed below in section IV B.

Kornreich and Scalo (1999) have recently considered the problem of turbulence maintenance in molecular clouds, comparing the average time between external shock wave passages through a cloud with the energy decay time, finding that they are comparable, and implying that this “shock pump” is capable of sustaining the cloud turbulence. An additional result is the proposal of a cascadelike mechanism for the compressible case, in which vorticity is generated behind shocks, which in turn rapidly produces new smaller-scale shocks, and so on. This is due to an interesting asymmetry of the evolution equations for the vorticity and divergence of the velocity field (for recent discussions, see Vázquez-Semadeni et al. 1996; Kornreich and Scalo 1999): the nonlinear transfer will produce compressible modes out of purely rotational ones, but the converse is not true. This can also be understood in terms of the well-known Kelvin theorem of conservation of circulation.

The mechanisms of vorticity production have also been investigated numerically. Simulations of compressible turbulence with purely potential forcing indicate that a negligible amount of kinetic energy is transferred from compressible to solenoidal modes (Kida and Orszag 1990; Vázquez-Semadeni et al. 1996). Vorticity generation behind curved shocks or shock intersections, as well as the vortex stretching term, do not seem to be efficient processes to maintain a nonnegligible level of vorticity in the flow. However, in the presence of the Coriolis force, thermal heating (through the baroclinic term), or magnetic field, near equipartition

between solenoidal and compressible modes is easily obtained (Vázquez-Semadeni et al. 1996).

## B. Turbulence and Cloud Support against Gravity

An issue that is often discussed in tandem with turbulent dissipation is the question of cloud support against self-gravity. Cold, dark clouds and GMCs have typical Jeans lengths  $\sim 2$  pc, such that the whole cloud entities exceed the Jeans mass  $M_J = \rho L_J^3$  by factors of a thousand or much more. This is just another way of stating that thermal pressure gradients would be powerless to prevent self-gravitating runaway. If not for the intervention of other dynamical processes, wholesale cloud collapse would be the rule.

“Turbulent pressure” is often invoked as a means to counter gravity. For a weakly compressible medium, the  $\rho v^2$  Reynolds stresses are naturally associated with this turbulent pressure. The effect of turbulence on the gravitational instability in the absence of a magnetic field was first investigated by Chandrasekhar (1951), who suggested an increase of the Jeans length, and later by Bonazzola et al. (1987, 1992) and Vázquez-Semadeni and Gazol (1995), who predicted a reversal of the Jeans criterion when certain conditions on the energy spectrum are met.

In a strongly compressible and radiative medium, the collisions of supersonically converging streams of gas are nearly inelastic, so the Reynolds stress does not itself act as an effective pressure. Turbulent motions can, however, generate turbulent magnetic fields (and vice versa); these fluctuating magnetic fields exert pressure and tension forces on the medium. Several authors (Shu et al. 1987; Fazio and Adams 1993; McKee and Zweibel 1995) have suggested that, in particular, the time-dependent magnetic field perturbations associated with Alfvén waves could potentially be important in providing “wave pressure” support against gravity along the mean magnetic field direction in a cloud. This mean-field axis is the most susceptible to collapse; in the orthogonal directions, magnetic pressure suppresses gravitational instability in a homogeneous cloud as long as  $n_J \equiv L/L_J < (\beta/2)^{-1/2}$  (Chandrasekhar and Fermi 1953) and suppresses instability in a cloud pancake of surface density  $\Sigma$  provided  $\Sigma/B < 1/(2\pi \sqrt{G})$  (cf. Mouschovias and Spitzer 1976; Tomisaka et al. 1988).

An exact derivation of the Jeans criterion is hardly possible in the MHD turbulent case. An attempt in this direction has been made by considering the linear stability of a self-gravitating medium, permeated by a uniform magnetic field  $B_0$  along which a finite-amplitude, circularly polarized Alfvén wave propagates. For perturbations along  $B_0$  it is found that the Alfvén wave increases the critical Jeans length (Lou 1997). In the case of perturbations perpendicular to the mean field, Gazol and Pasot (1999b) have shown that the medium is less stable in the presence of a moderate-amplitude Alfvén wave. For large-amplitude waves, however, McKee and Zweibel (1995) show that the waves have an isotropic stabilizing effect.

Gammie and Ostriker (1996) verified, using simulations in  $1\frac{2}{3}$ D, that Alfvén waves of sufficient amplitude (such that  $n_J \lesssim \delta v_A/2c_s$ ) can indeed prevent collapse of slab clouds along the mean field direction. These simulations included both cases with decaying and cases with quasisteady forced turbulence. For 1D decay models, the turbulent dissipation rate is low enough that clouds with initial turbulent energy above the limit remain uncollapsed for times up to  $t_g \equiv L_J/c$  ( $\sim 10$  Myr for typical conditions). However, more recent simulations performed in higher dimensions have shown that the greater dissipation rates quench turbulence too rapidly for magnetic fluctuations to prevent mean-field collapse. Because 3D dissipation times for both magnetized and unmagnetized flows are shorter than the flow-crossing time  $t_f$  (cf. section IV.A), they will also be less than the gravitational collapse times  $\approx 0.3t_g$  for clouds that are virialized (e.g., observed cloud scalings yield  $t_f \sim 0.5t_g$ ); thus, cloud support cannot be expected for unregenerated turbulence. Self-gravitating, magnetized  $2\frac{1}{2}$ D simulations of Ostriker et al. (1999) have already demonstrated this result directly and concluded that without ongoing energy inputs, only the strength of the mean magnetic field is important in determining whether or not clouds collapse in times  $< 10$  Myr. Numerical simulations have also been performed that allow for ongoing turbulent excitation. In unmagnetized 2D models, Léorat et al. (1990) showed that large-scale gravitational collapse can be prevented indefinitely provided a high enough Mach number is maintained by forcing, and the energy is injected at small enough scales. For magnetized models, Gammie and Ostriker (1996) found similar results for  $1\frac{2}{3}$ D simulations (as well as for unpublished 2D simulations).

Ballesteros-Paredes and Vázquez-Semadeni (1997) have measured the overall virial balance of clouds in a “survey” of the 2D high-resolution ISM simulations of Passot et al. (1995). It was found that the gravitational term appearing in the virial theorem is comparable to the sum of the other virial terms for the largest clouds but progressively loses importance on the average as smaller clouds are considered. Nevertheless, the scatter about this average trend is large, and a small fraction of small clouds have very large gravitational terms that overwhelm the others and induce collapse. In this scenario, the low efficiency of star formation is understood as a consequence of the intermittency of the turbulence.

### C. Evolution of the Star Formation Rate

Numerical simulations on the large scales provide information on the star formation (SF) history as well (Vázquez-Semadeni et al. 1995; Gazol and Passot 1998*a*). In these models, stars form whenever a certain density threshold is exceeded, rather than being treated as a separate fluid (e.g., Chiang and Prendergast 1985; Rosen et al. 1993). A nontrivial result is the development of a self-sustained cycle of SF in which the turbulence, aided by self-gravity, contains enough power to produce star-forming clouds, while the energy injected by the stars is sufficient to regenerate the turbulence. Due to the strong nonlinearity of the SF scheme, this cycle is

highly chaotic and particularly intermittent in the presence of supernovae. Self-propagating SF in supershells is very efficient, increasing the SF rate in active periods, but the destructive power of superbubbles is so large that the system requires a longer time to create new SF sites once the shells have dispersed. A similar result was obtained by Vázquez and Scalo (1989) in a simple model for gas-accreting galaxies. This behavior is consistent with recent observations suggesting that the SF history in galaxies is highly irregular (e.g., Grebel 1998). Another interesting point concerns the influence of the strength of the uniform component of the magnetic field  $B_0$  (Gazol and Passot 1998a). The star formation rate is found to increase with  $B_0$  (with  $B_0 = 5 \mu\text{G}$  it is larger by a factor  $\sim 3$  compared with the case  $B_0 = 0$ ), as long as  $B_0$  is not too large. For very large  $B_0$ , the SF rate decreases due to the rigidification of the medium.

## V. CONCLUSIONS

In this chapter we have reviewed a vast body of results that address the problem of MHD compressible (MHDC) turbulence and its implications for problems of cloud and star formation. Most of these results are new, appearing after the last *Protostars and Planets* conference (PP II). Interstellar turbulence is inherently a multiscale and nonequilibrium phenomenon and thus appears to play a fundamental role in the formation, evolution, and determination of cloud structural properties. Most of the studies reviewed here have relied on direct numerical simulations of MHDC turbulence in a variety of regimes, ranging from isothermal to polytropic to fully thermodynamic, the latter including parameterized heating and cooling and modeled star formation. High-resolution 2D and 3D simulations of the ISM at “box” scales from 1 kpc down to a few pc have shown that the clouds formed in them reproduce several observational cloud properties well (such as clumpy structure and line-width-size scalings), while suggesting that some others [e.g., Larson’s (1981) density-size relation] may either arise from selection effects or require other special conditions. Simulations to date have shown that the old paradigm that MHDC turbulence should dissipate much more slowly than nonmagnetic turbulence may be incorrect, bringing back the necessity of strong large-scale fields for magnetic cloud support and of continued energy injection in order to sustain the turbulence. The idea for self-regulation of star formation by turbulent feedback remains promising, and it is also possible that physical processes not yet incorporated in simulations may reduce the turbulent dissipation rate.

A large number of questions remain unanswered, however. At the larger scales it is necessary to understand the interplay between turbulence, large-scale instabilities, and spiral waves in the formation of molecular clouds and complexes, as well as to investigate the processes that lead to cloud destruction. Quantitative assessments must be developed for the efficiency of turbulent excitation from internal and external sources. A

theory of the IMF, or at least of the cloud mass spectrum, requires the knowledge and understanding of multipoint density statistics. Better understanding of the parameter dependence of these and other questions in cloud evolution and structure will be required to address the problem of “deriving” the star formation rate and efficiency, in answer to the oft-posed challenge of extragalactic astronomers. Comparison with observations is crucial to discriminate among models, but it will always face the degeneracy limitations associated with projection effects.

Future research in these areas is likely to include more physical processes (e.g., ambipolar diffusion, radiative transfer, and global environment effects, such as the galactic spiral potential and variations in external radiation with galactic position), to perform direct statistical comparison with observations of molecular lines, and polarization and IR measurements, and to work at higher resolutions in 3D, if the multiscale nature of the problem is to be captured adequately. With so much still untried, there is considerable ground to cover before reaching the long-range goal of synthesizing the results of disparate numerical simulations into a coherent theory of the turbulent ISM. Nevertheless, the important advances since PP III show the success of numerical methods in answering many long-standing questions in the theory of interstellar MHD turbulence and the opportunity for great progress before the next *Protostars and Planets*.

*Acknowledgments* We are grateful to C. F. McKee for helpful comments on this manuscript and to J. Scalo for enlightening discussions. This work has been supported in part by grants UNAM/DGAPA IN105295, CRAY/UNAM SC008397, NAG13380 from NASA, and by the CNRS “PCMI” National Program.

## REFERENCES

- Arons, J., and Max, C. E. 1975. Hydromagnetic waves in molecular clouds. *Astrophys. J. Lett.* 196:77–81
- Ballesteros-Paredes, J., and Vázquez-Semadeni, E. 1997. Virial balance and scaling relations in turbulent MHD two-dimensional numerical simulations of the ISM. In *Star Formation Near and Far*, ed. S. Holt and L. Mundy (Woodbury, NY: AIP Press), pp. 81–84.
- Ballesteros-Paredes, J., Vázquez-Semadeni, E., and Scalo, J. 1999. Clouds as turbulent density fluctuations. Implications for pressure confinement and spectral line data interpretation. *Astrophys. J.* 515:286–303.
- Balsara, D. S., Crutcher, R. M., and Pouquet, A. 1997. Numerical simulations of MHD turbulence and gravitational collapse. In *Star Formation, Near and Far*, ed. S. Holt and L. Mundy (Woodbury, NY: AIP Press), pp. 89–92.
- Bash, F. N., Green, E., and Peters, W. L., III. 1977. The galactic density wave, molecular clouds, and star formation. *Astrophys. J.* 217:464–472.
- Bate, M. R., Clarke, C. J., and McCaughrean, M. J. 1998. Interpreting the mean surface density of companions in star-forming regions. *Mon. Not. Roy. Astron. Soc.* 297:1163–1181.

- Bertoldi, F., and McKee, C. 1992. Pressure-confined clumps in magnetized molecular clouds. *Astrophys. J.* 395:140–157.
- Blitz, L. 1993. Giant molecular clouds. In *Protostars and Planets III*, ed. E. H. Levy and J. I. Lunine (Tucson: University of Arizona Press), pp. 125–161.
- Blitz, L. 1994. Simple things we don't know about molecular clouds. In *The Cold Universe*, ed. T. Montmerle, C. J. Lada, I. F. Mirabel, and J. Trân Thanh Vân (Gif-sur-Yvette: Editions Frontières), pp. 99–106.
- Blitz, L., and Shu, F. H. 1980. The origin and lifetime of giant molecular cloud complexes. *Astrophys. J.* 238:148–157.
- Blitz, L., and Williams, J. P. 1997. Molecular clouds are not fractal: A characteristic size scale in Taurus. *Astrophys. J. Lett.* 488:L145–L148.
- Bonazzola, S., Falgarone, E., Heyvaerts, J., Péroul, M., and Puget, J. L. 1987. Jeans collapse in a turbulent medium. *Astron. Astrophys.* 172:293–298.
- Bonazzola, S., Péroul, M., Puget, J. L., Heyvaerts, J., Falgarone, E., and Panis, J. F. 1992. Jeans collapse of turbulent gas clouds. A tentative theory. *J. Fluid Mech.* 245:1–28.
- Boulares, A., and Cox, D. P. 1990. Galactic hydrostatic equilibrium with magnetic tension and cosmic-ray diffusion. *Astrophys. J.* 365:544–558.
- Braun, R. 1999. Properties of atomic gas in spiral galaxies. In *Interstellar Turbulence, Proceedings of the 2nd Guillermo Haro Conference*, ed. J. Franco and A. Carramiñana (Cambridge: Cambridge University Press), pp. 12–19.
- Burgers, J. M. 1974. *The Nonlinear Diffusion Equation* (Dordrecht: Reidel).
- Carr, J. S. 1987. A study of clumping in the Cepheus OB 3 molecular cloud. *Astrophys. J.* 323:170–178.
- Chandrasekhar, S. 1951. The gravitational instability in an infinite homogeneous turbulent medium. *Proc. Royal Soc. London* 210:26–29.
- Chandrasekhar, S. 1961. *Hydrodynamic and Hydromagnetic Stability* (Oxford: Clarendon Press).
- Chandrasekhar, S., and Fermi, E. 1953. Problems of gravitational instability in the presence of a magnetic field. *Astrophys. J.* 118:116–141.
- Chappell, D., and Scalo, J. 1998. Multifractal scaling, geometrical diversity, and hierarchical structure in the cool interstellar medium. *Astrophys. J.*, in press.
- Chiang, W.-H., and Prendergast, K. H. 1985. Numerical study of a two-fluid hydrodynamic model of the interstellar medium and Population I stars. *Astrophys. J.* 297:507–530.
- Crutcher, R. M. 1999. Observations of magnetic fields in dense clouds: Implications for MHD turbulence and cloud evolution. In *Interstellar Turbulence, Proceedings of the 2nd Guillermo Haro Conference*, ed. J. Franco and A. Carramiñana (Cambridge: Cambridge University Press), pp. 213–217.
- Crutcher, R. M., Troland, T. H., Goodman, A. A., Heiles, C., Kazès, I., and Myers, P. C. 1993. OH Zeeman observations of dark clouds. *Astrophys. J.* 407:175–184.
- Crutcher, R. M., Troland, T. H., Lazareff, B., Kazès, I. 1996. CN Zeeman observations of molecular cloud cores. *Astrophys. J.* 456:217–224.
- de Jong, T., Dalgarno, A., and Boland, W. 1980. Hydrostatic models of molecular clouds. I. Steady state models. *Astron. Astrophys.* 91:68–84.
- Draine, B. T. 1978. Photoelectric heating of interstellar gas. *Astrophys. J. Suppl.* 36:595–619.
- Elmegreen, B. G. 1991. Cloud formation by combined instabilities in galactic gas layers. Evidence for a Q threshold in the fragmentation of shearing wavelets. *Astrophys. J.* 378:139–156.
- Elmegreen, B. G. 1993a. Formation of interstellar clouds and structure. In *Protostars and Planets III*, ed. E. H. Levy and J. I. Lunine (Tucson: University of Arizona Press), pp. 97–124.

- Elmegreen, B. G. 1993*b*. Star formation at compressed interfaces in turbulent self-gravitating clouds. *Astrophys. J. Lett.* 419:L29–L32.
- Elmegreen, B. G. 1995. Star formation on a large scale. In *Molecular Clouds and Star Formation*, ed. C. Yuan and Junhan You (Singapore: World Scientific), pp. 149–205.
- Elmegreen, B. G., and Lada, C. J. 1977. Sequential formation of subgroups in OB associations. *Astrophys. J.* 214:725–741.
- Elmegreen, B. G., and Elmegreen, D. M. 1978. Star formation in shock-compressed layers. *Astrophys. J.* 220:1051–1062.
- Falgarone, E., Phillips, T. G., and Walker, C. K. 1991. The edges of molecular clouds: Fractal boundaries and density structure. *Astrophys. J.* 378:186–200.
- Falgarone, E., Puget, J.-L., and Pérault, M. 1992. The small-scale density and velocity structure of molecular clouds. *Astron. Astrophys.* 257:715–730.
- Falgarone, E., Panis, J.-F., Heithausen, A., Pérault, M., Stutzki, J., Puget, J.-L., and Bensch, F. 1998. The IRAM key-project: Small-scale structure of pre-star-forming regions. I. Observational results. *Astron. Astrophys.* 331:669–696.
- Fatuzzo, M., and Adams, F. C. 1993. Magnetohydrodynamic wave propagation in one-dimensional nonhomogeneous, self-gravitating clouds. *Astrophys. J.* 412:146–159.
- Ferrini, F., Marchesoni, F., and Vulpiani, A. 1983. A hierarchical model for gravitational compressible turbulence. *Astrophys. Space Sci.* 96:83–93.
- Field, G. B., Goldsmith, D. W., and Habing, H. J. 1969. Cosmic ray heating of the interstellar gas. *Astrophys. J. Lett.* 155:L149–L154.
- Fleck, R. C. 1981. On the generation and maintenance of turbulence in the interstellar medium. *Astrophys. J. Lett.* 246:L151–L154.
- Fleck, R. C. 1996. Scaling relations for the turbulent, non-self-gravitating, neutral component of the interstellar medium. *Astrophys. J.* 458:739–741.
- Gammie, C. F. 1996. Linear theory of magnetized, viscous, self-gravitating gas disks. *Astrophys. J.* 462:725–731.
- Gammie, C. F., and Ostriker, E. C. 1996. Can nonlinear hydromagnetic waves support a self-gravitating cloud? *Astrophys. J.* 466:814–830.
- Gazol, A., and Passot, T. 1999*a*. A turbulent model for the interstellar medium. III. Stratification and supernova explosions. *Astrophys. J.* 518:748–759.
- Gazol, A., and Passot, T. 1999*b*. Influence of Alfvén waves on the gravitational instability. *Astron. Astrophys.*, submitted.
- Goldreich, P., and Kwan, J. 1974. Molecular clouds. *Astrophys. J.* 189:441–453.
- Goldreich, P., and Sridhar, S. 1995. Toward a theory of interstellar turbulence. II: Strong Alfvénic turbulence. *Astrophys. J.* 438:763–775.
- Goodman, A. A., Barranco, J. A., Wilner, D. J., and Heyer, M. H. 1998. Coherence in dense cores. II. The transition to coherence. *Astrophys. J.* 504:223–246.
- Grebel, E. 1998. Star formation histories of Local Group dwarf galaxies. In *Dwarf Galaxies and Cosmology*, ed. T. X. Thuan, C. Balkowski, V. Cayatte, and J. Trân Thanh Vân (Gif-sur-Yvette: Editions Frontières), in press.
- Heiles, C., Goodman, A. A., McKee, C. F., and Zweibel, E. G. 1993. Magnetic fields in star-forming regions: Observations. In *Protostars and Planets III*, ed. E. H. Levy and J. I. Lunine (Tucson: University of Arizona Press), pp. 279–326.
- Henriksen, R. N., and Turner, B. E. 1984. Star cloud turbulence. *Astrophys. J.* 287:200–297.
- Heyer, M. H., Carpenter, J. M., and Ladd, E. F. 1996. Giant molecular cloud complexes with optical H II regions:  $^{12}\text{CO}$  and  $^{13}\text{CO}$  observations and global cloud properties. *Astrophys. J.* 463:630–641.
- Houlahan, P., and Scalo, J. 1990. Recognition and characterization of hierarchical interstellar structure. I. Correlation function. *Astrophys. J. Suppl.* 72:133–152.

- Houlahan, P., and Scalo, J. 1992. Recognition and characterization of hierarchical interstellar structure. II. Structure tree statistics. *Astrophys. J.* 393:172–187.
- Hunter, J. H., Jr. 1979. The influence of initial velocity fields upon star formation. *Astrophys. J.* 233:946–949.
- Hunter, J. H., Jr., and Fleck, R. C. 1982. Star formation: The influence of velocity fields and turbulence. *Astrophys. J.* 256:505–513.
- Hunter, J. H., Jr., Sandford, M. T., II, Whitaker, R. W., and Klein, R. I. 1986. Star formation in colliding gas flows. *Astrophys. J.* 305:309–332.
- Issa, M., MacLaren, I., and Wolfendale, W. 1990. The size–line width relation and the mass of molecular hydrogen. *Astrophys. J.* 352:132–138.
- Kadomtsev, B. B., and Petviashvili, V. I. 1973. Acoustic turbulence. *Sov. Phys. Dokl.* 18:115–116.
- Kegel, W. H. 1989. The interpretation of correlation between observed parameters of molecular clouds. *Astron. Astrophys.* 225:517–520.
- Kida, S., and Orszag, S. A. 1990. Energy and spectral dynamics in forced compressible turbulence. *J. Sci. Comput.* 5:85–125.
- Klein, R. I., and Woods, D. T. 1998. Bending mode instabilities and fragmentation in interstellar cloud collisions: A mechanism for complex structure. *Astrophys. J.* 497:777–799.
- Kolmogorov, A. N. 1941. The local structure of turbulence in an incompressible viscous fluid for very large Reynolds numbers. *Dokl. Akad. Nauk* 30:301–305.
- Kornreich, P., and Scalo, J. 1999. The galactic shock pump. A source of supersonic internal motions in the cool interstellar medium. *Astrophys. J.*, in press.
- Kraichnan, R. H. 1968. Lagrangian-history statistical theory for Burgers' equation. *Phys. Fluids* 11:265–277.
- Kulsrud, R. M., and Pearce, W. P. 1969. The effect of wave-particle interactions on the propagation of cosmic rays. *Astrophys. J.* 163:567–576.
- Lada, E. A., Evans, N. J., II, and Falgarone, E. 1997. Physical properties of molecular cloud cores in L1630 and implications for star formation. *Astrophys. J.* 488:286–306.
- Landau, L. D., and Lifshitz, E. M. 1987. *Fluid Mechanics*. 2nd ed. (Oxford: Pergamon Press).
- Larson, R. B. 1981. Turbulence and star formation in molecular clouds. *Mon. Not. Roy. Astron. Soc.* 194:809–826.
- Larson, R. B. 1995. Star formation in groups. *Mon. Not. Roy. Astron. Soc.* 272:213–220.
- Lazarian, A. 1995. Study of turbulence in H I using radiointerferometers. *Astron. Astrophys.* 293:507–520.
- Léorat, J., Passot, T., and Pouquet, A. 1990. Influence of supersonic turbulence on self-gravitating flows. *Mon. Not. Roy. Astron. Soc.* 243:293–311.
- Loren, R. B. 1989. The cobwebs of Ophiuchus. I. Strands of (C-13)O: The mass distribution. *Astrophys. J.* 338:902–924.
- Lou, Y.-Q. 1997. Gravitational collapse in the presence of a finite-amplitude circularly polarized Alfvén wave. *Mon. Not. Roy. Astron. Soc.* 279:L67–L71.
- Mac Low, M.-M. 1999. The energy dissipation rate of supersonic, magnetohydrodynamic turbulence in molecular clouds. *Astrophys. J.*, submitted.
- Mac Low, M.-M., Klessen, R. S. Burkert, and Smith, M. D. 1998. Kinetic energy decay of supersonic and super-Alfvénic turbulence in star-forming clouds. *Phys. Rev. Lett.* 80:2754–2757.
- Maloney, P. 1988. The turbulent interstellar medium and pressure-bounded molecular clouds. *Astrophys. J.* 334:761–770.
- McKee, C. F. 1989. Photoionization-regulated star formation and the structure of molecular clouds. *Astrophys. J.* 345:782–801.

DR. P. RUPAK NATH

- McKee, C. F., and Zweibel, E. G. 1992. On the virial theorem for turbulent molecular clouds. *Astrophys. J.* 399:551–562.
- McKee, C. F., and Zweibel, E. G. 1995. Alfvén waves in interstellar gas dynamics. *Astrophys. J.* 440:686–696.
- McKee, C. F., Zweibel, E. G., Goodman, A. A., and Heiles, C. 1993. Magnetic fields in star-forming regions. In *Protostars and Planets III*, ed. E. H. Levy and J. I. Lunine (Tucson: University of Arizona Press), pp. 327–366.
- Miesch, M. S., and Bally, J. 1994. Statistical analysis of turbulence in molecular clouds. *Astrophys. J.* 429:645–671.
- Mouschovias, T. 1987. Star formation in magnetic interstellar clouds. I. Interplay between theory and observation. In *Physical Processes in Interstellar Clouds*, ed. G. E. Morfill and M. Scholer (Dordrecht: Reidel), pp. 453–489.
- Mouschovias, T., and Spitzer, L. 1976. Note on the collapse of magnetic interstellar clouds. *Astrophys. J.* 210:326–327.
- Myers, P. C. 1978. A compilation of interstellar gas properties. *Astrophys. J.* 225:380–389.
- Myers, P. C. 1995. Star forming molecular clouds. In *Molecular Clouds and Star Formation*, ed. C. Yuan and Junhan You (Singapore: World Scientific), pp. 47–96.
- Myers, P. C., and Goodman, A. A. 1988a. Evidence for magnetic and virial equilibrium in molecular clouds. *Astrophys. J. Lett.* 326:27–30.
- Myers, P. C., and Goodman, A. A. 1988b. Magnetic molecular clouds: Indirect evidence for magnetic support and ambipolar diffusion. *Astrophys. J.* 329:392–405.
- Myers, P. C., and Khersonsky, V. K. 1995. On magnetic turbulence in interstellar clouds. *Astrophys. J.* 442:186–196.
- Nordlund, Å., and Padoan, P. 1999. The density PDFs of supersonic random flows. In *Interstellar Turbulence, Proceedings of the 2nd Guillermo Haro Conference*, ed. J. Franco and A. Carramiñana (Cambridge: Cambridge University Press), pp. 218–222.
- Norman, C., and Ferrara, A. 1996. The turbulent interstellar medium: Generalizing to a scale-dependent phase continuum. *Astrophys. J.* 467:280–291.
- Norman, C., and Silk, J. 1980. Clumpy molecular clouds: A dynamic model self-consistently regulated by T Tauri star formation. *Astrophys. J.* 238:158–174.
- Obukhov, A. M. 1941. Of the distribution of energy in the spectrum of turbulent flow. *Comptes Rend. Acad. Sci. URSS* 32:19.
- Ögelman, H. B., and Maran, S. P. 1976. The origin of OB associations and extended regions of high-energy activity in the Galaxy through supernova cascade processes. *Astrophys. J.* 209:124–129.
- Ostriker, E. C. 1997. Turbulence and magnetic fields in star formation. In *Star Formation, Near and Far*, ed. S. Holt and L. Mundy (Woodbury, NY: AIP Press), pp. 51–63.
- Ostriker, E. C., Gammie, C. F., and Stone, J. M. 1999. Kinetic and structural evolution of self-gravitating, magnetized clouds: 2.5-dimensional simulations of decaying turbulence. *Astrophys. J.* 513:259–274.
- Padoan, P. 1995. Supersonic turbulent flows and the fragmentation of a cold medium. *Mon. Not. Roy. Astron. Soc.* 277:377–388.
- Padoan, P., and Nordlund, Å. 1999. A super-Alfvénic model for dark clouds. *Astrophys. J.*, in press.
- Padoan, P., Jones, B. J. T., Nordlund, Å. 1997a. Supersonic turbulence in the interstellar medium: Stellar extinction determinations as probes of the structure and dynamics of dark clouds. *Astrophys. J.* 474:730–734.
- Padoan, P., Nordlund, Å., and Jones, B. J. T. 1997b. The universality of the stellar initial mass function. *Mon. Not. Roy. Astron. Soc.* 288:145–152.

- Padoan, P., Juvela, M., Bally, J., and Nordlund, Å. 1998. Synthetic molecular clouds from supersonic MHD and non-LTE radiative transfer calculations. *Astrophys. J.* 504:300–313.
- Passot, T., and Vázquez-Semadeni, E. 1998. Density probability distribution in one-dimensional gas dynamics. *Phys. Rev. E* 58:4501–4510.
- Passot, T., Pouquet, A., and Woodward, P. 1988. The plausibility of Kolmogorov-type spectra in molecular clouds. *Astron. Astrophys.* 197:228–234.
- Passot, T., Vázquez-Semadeni, E., and Pouquet, A. 1995. A turbulent model for the interstellar medium. II. Magnetic fields and rotation. *Astrophys. J.* 455:536–555.
- Peng, R., Langer, W. D., Velusamy, T., Kuiper, T. B. H., and Levin, S. 1998. Low-mass clumps in TMC-1: Scaling laws in the small-scale regime. *Astrophys. J.* 197:842–849.
- Peters, W. L., and Bash, F. N. 1987. The correlation of spiral arm molecular clouds with atomic hydrogen. *Astrophys. J.* 317:646–652.
- Plume, R., Jaffe, D. T., Evans, N. J., II, Martín-Pintado, J., and Gómez-González, J. 1997. Dense gas and star formation: Characteristics of cloud cores associated with water masers. *Astrophys. J.* 476:730–749.
- Pouquet, A., Passot, T., and Léorat, J. 1990. Numerical simulations of compressible flows. Proceedings of the IAU symposium 147. In *Fragmentation of Molecular Clouds and Star Formation*, ed. E. Falgarone, F. Boulanger, and G. Duvert (Dordrecht: Kluwer), pp. 101–118.
- Pratap, P., Dickens, J. E., Snell, R. L., Miralles, M. P., Bergin, E. A., Irvine, W. M., and Schloerb, F. P. 1997. A study of the physics and chemistry of TMC-1. *Astrophys. J.* 486:862–885.
- Rosen, A., Bregman, J. N., and Norman, M. L. 1993. Hydrodynamical simulations of star-gas interactions in the interstellar medium with an external gravitational potential. *Astrophys. J.* 413:137–149.
- Scalo, J. 1985. Fragmentation and hierarchical structure in the interstellar medium. In *Protostars and Planets II*, ed. D. C. Black and M. S. Matthews (Tucson: University of Arizona Press), pp. 201–296.
- Scalo, J. 1987. Theoretical approaches to interstellar turbulence. In *Interstellar Processes*, ed. D. J. Hollenbach and H. A. Thompson, Jr. (Dordrecht: Reidel), pp. 349–392.
- Scalo, J. 1990. Perception of interstellar structure. Facing complexity. In *Physical Processes in Fragmentation and Star Formation*, ed. R. Capuzzo-Dolcetta, C. Chiosi, and A. di Fazio (Dordrecht: Kluwer), pp. 151–177.
- Scalo, J., Vázquez-Semadeni, E., Chappell, D., and Passot, T. 1998. On the probability density function of galactic gas. I. Numerical simulations and the significance of the polytropic index. *Astrophys. J.* 504:835–853.
- Shaya, E. J., and Federman, S. R. 1987. The H I distribution in clouds within galaxies. *Astrophys. J.* 319:76–83.
- Shu, F. H., Adams, F. C., and Lizano, S. 1987. Star formation in molecular clouds: Observation and theory. *Ann. Rev. Astron. Astrophys.* 25:23–81.
- Simon, M. 1997. Clustering of young stars in Taurus, Ophiuchus, and the Orion Trapezium. *Astrophys. J. Lett.* 482:L81–L84.
- Solomon, P. M., and Sanders, D. B. 1985. Star formation in a galactic context: The location and properties of molecular clouds. In *Protostars and Planets II*, ed. D. C. Black and M. S. Matthews (Tucson: University of Arizona Press), pp. 59–80.
- Spitzer, L., Jr., and Savedoff, M. P. 1950. The temperature of interstellar matter. III. *Astrophys. J.* 111:593–608.
- Stevens, I. R., Blondin, J. M., and Pollock, A. M. T. 1992. Colliding winds from early-type stars in binary systems. *Astrophys. J.* 386:265–287.

- Stone, J. M. 1999. Direct numerical simulations of compressible MHD turbulence. In *Interstellar Turbulence, Proceedings of the 2nd Guillermo Haro Conference*, ed. J. Franco and A. Carramiñana (Cambridge: Cambridge University Press), pp. 267–271.
- Stone, J. M., Ostriker, E. C., and Gammie, C. F. 1998. Dissipation in compressible MHD turbulence. *Astrophys. J. Lett.* 508:L99–L102.
- Tohline, J. E., Bodenheimer, P. H., and Christodoulou, D. M. 1987. The crucial role of cooling in the making of molecular clouds and stars. *Astrophys. J.* 322:787–794.
- Tomisaka, K., Ikeuchi, S., and Nakamura T. 1988. Equilibria and evolutions of magnetized, rotating, isothermal clouds. II. The extreme case: Nonrotating clouds. *Astrophys. J.* 335:239–262.
- Torrelles, J. M., Rodríguez, L. F., Cantó, J., Marcaide, J., and Gyulbudagian, A. L. 1983. A search for molecular outflows associated with peculiar nebulosities and regions of star formation. *Rev. Mex. Astron. Astrofís.* 8:147–154.
- Troland, T. H., Crutcher, R. M., Goodman, A. A., Heiles, C., Kamez, L., and Myers, P. C. 1996. The magnetic fields in the Ophiuchus and Taurus molecular clouds. *Astrophys. J.* 471:302–307.
- Vázquez, E. C., and Scalo, J. 1989. Evolution of the star formation rate in galaxies with increasing densities. *Astrophys. J.* 343:644–658.
- Vázquez-Semadeni, E. 1994. Hierarchical structure in nearly pressureless flows as a consequence of self-similar statistics. *Astrophys. J.* 423:681–692.
- Vázquez-Semadeni, E. 1999. Turbulence in molecular clouds. In *Millimeter and Submillimeter Astronomy: Chemistry and Physics in Molecular Clouds*, ed. W. F. Wall, A. Carramiñana, L. Carrasco, and P. F. Goldsmith (Dordrecht: Kluwer), pp. 161–186.
- Vázquez-Semadeni, E., and Gazol, A. 1995. Gravitational instability in turbulent, non-uniform media. *Astron. Astrophys.* 303:204–210.
- Vázquez-Semadeni, E., and Passot, T. 1999. Turbulence as an organizing agent in the ISM. In *Interstellar Turbulence, Proceedings of the 2nd Guillermo Haro Conference*, ed. J. Franco and A. Carramiñana (Cambridge: Cambridge University Press), pp. 223–231.
- Vázquez-Semadeni, E., Passot, T., and Pouquet, A. 1995. A turbulent model for the interstellar medium. I. Threshold star formation and self-gravity. *Astrophys. J.* 441:702–725.
- Vázquez-Semadeni, E., Passot, T., and Pouquet, A. 1996. Influence of cooling-induced compressibility on the structure of turbulent flows and gravitational collapse. *Astrophys. J.* 473:881–893.
- Vázquez-Semadeni, E., Ballesteros-Paredes, J., and Rodríguez, L. F. 1997. A search for Larson-type relations in numerical simulations of the ISM. Evidence for non-constant column densities. *Astrophys. J.* 474:292–307.
- Vázquez-Semadeni, E., Cantó, J., and Lizano, S. 1998. Does turbulent pressure behave as a logatope? *Astrophys. J.* 492:596–602.
- Vishniac, E. T. 1983. The dynamic and gravitational instabilities of spherical shocks. *Astrophys. J.* 274:152–167.
- Vishniac, E. T. 1994. Nonlinear instabilities in shock-bounded slabs. *Astrophys. J.* 428:186–208.
- Wannier, P. G., Lichten, S. M., Andersson, B.-G., and Morris, M. 1991. Warm neutral halos around molecular clouds. II. HI and CO ( $J = 1-0$ ) observations. *Astrophys. J. Suppl.* 75:987–998.
- Whitworth, A. 1996. The structure of the neutral interstellar medium: A theory of interstellar turbulence. In *Unsolved Problems of the Milky Way*, ed. L. Blitz and P. Teuben (Dordrecht:Kluwer) pp. 591–596.

- Williams, J. P., Blitz, L., and Stark, A. A. 1995. The density structure in the Rosette molecular cloud: Signposts of evolution. *Astrophys. J.* 451: 252–274.
- Wilson, T. L., and Walmsley, C. M. 1989. Small-scale clumping in molecular clouds. *Astron. Astrophys. Rev.* 1:141-176.
- Wolfire, M. G., Hollenbach, D., McKee, C. F., Tielens, A. G. G. M., and Bakes, E. L. O. 1995. The neutral atomic phases of the interstellar medium. *Astrophys. J.*, 443:152–168.

DR.RUPNATHJIK(DR.RUPAK NATH)